

Conformal radius: At the interface of traditions

Kazantsev A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Pleiades Publishing, Ltd. H. Behnke's and E. Peschl's definition of plänarkonvexität leads to the Epstein-type inequalities when applied to the Hartogs domains in \mathbb{C}^2 . One-parameter series of such inequalities reveals the following rigidity phenomenon: the set of the parameters with extensive inequalities is exactly the segment which center corresponds to the well-known Nehari ball. The latter plays the crucial role in the forming the Gakhov class of all holomorphic and locally univalent functions in the unit disk with no more than one-pointed null-sets of the gradients of their conformal radii. The sufficient condition for the piercing of the Nehari sphere out of the Gakhov class is found. We deduce such a condition along the lines of the subordination approach to the proof of Haegi's theorem about the inclusion of any convex holomorphic function into the Gakhov class.

<http://dx.doi.org/10.1134/S1995080217030167>

Keywords

Conformal radius, Epstein inequality, Gakhov class, Gakhov equation, Hartogs domain, hyperbolic derivative, linear convexity, linear-invariant family

References

- [1] H. Behnke and E. Peschl, "Zur Theorie der Funktionen mehrerer komplexer Veränderlichen. Konvexität in bezug auf analytische Ebenen im kleinen und großen," *Math. Ann.* 111, 158–177 (1935).
- [2] E. Peschl, "Über die Krümmung von Niveaulinien bei der konformen Abbildung einfach zusammenhängender Gebiete auf das Innere eines Kreises. Eine Verallgemeinerung eines Satzes von E. Study," *Math. Ann.* 106, 574–594 (1932).
- [3] E. Peschl, "Über die Verwendung von Differentialinvarianten bei gewissen Funktionenfamilien und die Übertragung einer darauf gegründeten Methode auf partielle Differentialgleichungen vom elliptischen Typus," *Ann. Acad. Sci. Fenn. Ser. A1: Math.* 336 (6), 1–22 (1963).
- [4] C. L. Epstein, "The hyperbolic Gauss map and quasiconformal reflections," *J. Reine Angew. Math.* 372, 96–135 (1986).
- [5] M. Chuaqui, "A unified approach to univalence criteria in the unit disk," *Proc. Am. Math. Soc.* 123, 441–453 (1995).
- [6] Ch. Pommerenke, "Linear-invariante Familien analytischer Funktionen," *Math. Ann.* 155, 108–154 (1964).
- [7] D. M. Campbell, "Locally univalent functions with locally univalent derivatives," *Trans. Am. Math. Soc.* 162, 395–409 (1971).
- [8] A. V. Kazantsev, "Bifurcations and new uniqueness criteria for critical points of hyperbolic derivatives," *Lob. J. Math.* 32, 426–437 (2011).
- [9] F. D. Gakhov, "On the inverse boundary problems," *Dokl. Akad. Nauk SSSR* 86, 649–652 (1952).
- [10] L. A. Aksent'ev, "The connection of the exterior inverse boundary value problem with the inner radius of the domain," *Izv. Vyssh. Uchebn. Zaved., Mat.* 2, 3–11 (1984).

- [11] L. A. Aksent'ev, Yu. E. Khokhlov, and E. A. Shirokova, "Uniqueness of solution of the exterior inverse boundary-value problem," *Mat. Zam.* 24, 672-678 (1978).
- [12] Yu. E. Khokhlov, "On the solvability of the exterior inverse boundary value problems for the analytic functions," *Dokl. Akad. Nauk SSSR* 278, 298-301 (1984).
- [13] S. R. Nasyrov and Yu. E. Khokhlov, "Uniqueness of the solution of the exterior inverse boundary value problem in the class of spiraloid domains," *Izv. Vyssh. Uchebn. Zaved., Mat.* 8, 24-27 (1984).
- [14] L. A. Aksent'ev, M. I. Kinder, and S. B. Sagitova, "Solvability of the exterior inverse boundary value problem in the case of multiply connected domain," *Tr. Semin. Kraev. Zadacham* 20, 22-34 (1983).
- [15] F. F. Mayer and M. A. Sevodin, "Conditions for univalence in domains with convex completion," *Tr. Semin. Kraev. Zadacham* 22, 151-160 (1985).
- [16] G. Polya and G. Szegő, *Problems and Theorems in Analysis* (Springer, New York, 1972; Nauka, Moscow, 1978), Vol.2.
- [17] H. R. Haegi, "Extremalprobleme und Ungleichungen konformer Gebietsgrößen," *Compos.Math.* 8, 81-111 (1950).
- [18] A. V. Kazantsev, "On a problem of Polya and Szegő," *Lob. J. Math.* 9, 37-46 (2001).
- [19] S. M. Einstein-Matthews, "Boundary behaviour of extremal plurisubharmonic functions," *Nagoya Math. J.* 138, 65-112 (1995).
- [20] S. V. Znamenskii, "Seven problems on C-convexity," in *Complex Analysis in Contemporary Mathematics, On the 80th Anniversary of the Birth of B. V. Shabat*, Ed. by E. M. Chirka (Fazis, Moscow, 2001), pp. 123-131 [in Russian].
- [21] A. V. Kazantsev, "Linear convexity of some Hartogs domain is Epsteinian," in *Proceedings of the 3rd International Conference on Geometric Analysis and its Applications, Volgograd, May 30-June 3, 2016* (Mat. Inst. SO RAN Soboleva, Volgograd Univ., Volgograd, 2016).
- [22] F. G. Avkhadiev, "Conditions for the univalence of analytic functions," *Izv. Vyssh. Uchebn. Zaved., Mat.* 11, 3-13 (1970).
- [23] J. A. Hummel, "The coefficient regions of starlike functions," *Pacif. J. Math.* 7, 1381-1389 (1957).
- [24] Z. Nehari, "The Schwarzian derivative and schlicht functions," *Bull. Am.Math. Soc.* 55, 545-551 (1949).
- [25] F. W. Gehring and Ch. Pommerenke, "On the Nehari univalence criterion and quasicircles," *Comm. Math. Helv.* 59, 226-242 (1984).
- [26] St. Ruscheweyh and K.-J. Wirths, "On extreme Bloch functions with prescribed critical points," *Math. Z.* 180, 91-106 (1982).
- [27] B. Kawohl, "Rearrangements and convexity of level sets in PDE," *Lect. NotesMath.* 1150, 1-136 (1985).
- [28] F. D. Gakhov, *Boundary Value Problems* (Fizmatlit, Moscow, 1958, 1963; Nauka, Moscow, 1977; Pergamon, Oxford, 1966).
- [29] L. A. Aksent'ev, A. V. Kazantsev, and A. V. Kiselev, "Uniqueness of the solution of an exterior inverse boundary value problem," *Izv. Vyssh. Uchebn. Zaved., Mat.* 10, 8-18 (1984).
- [30] L. A. Aksent'ev and A. V. Kazantsev, "A new property of the Nehari class and its application," *Izv. Vyssh. Uchebn. Zaved., Mat.* 8, 69-72 (1989).
- [31] G. G. Tumashev and M. T. Nuzhin, *Inverse Boundary Value Problems and Their Applications* (Kazan Univ., Kazan, 1965) [in Russian].
- [32] S. Yamashita, "The Schwarzian derivative and local maxima of the Bloch derivative," *Math. Jpn.* 37, 1117-1128 (1992).
- [33] M. Chuaqui and B. Osgood, "Ahlfors-Weill extensions of conformal mappings and critical points of the Poincaré metric," *Comment.Math. Helv.* 69, 659-668 (1994).
- [34] O. Martio and J. Sarvas, "Injectivity theorems in plane and space," *Ann. Acad. Sci. Fenn. Ser. A1: Math.* 4, 383-401 (1978-1979).
- [35] M. Chuaqui and B. Osgood, "Recent progress on the geometry of univalence criteria," *Contemp. Math.* 240, 75-87 (1999).
- [36] U. Eco, *La struttura assente: introduzione alla ricerca semiologia* (Bombiani, Milano, 1968; Petropolis, St. Petersburg, 1998) [in Italian].
- [37] A. V. Kazantsev, *Four Etudes on a Theme of F. D. Gakhov* (Marii-El Univ., Yoshkar-Ola, 2012) [in Russian].
- [38] R. Collins, *The Sociology of Philosophies. A Global Theory of Intellectual Change* (Belknap, Harvard Univ. Press, Cambridge, MA, London, 1998).
- [39] A. V. Kazantsev, "Bifurcations of roots of the Gakhov equation with a Löwner left-hand side," *Izv. Vyssh. Uchebn. Zaved., Mat.* 6, 69-73 (1993).